

Available online at [ScienceDirect](https://www.sciencedirect.com)

# Resuscitation

journal homepage: [www.elsevier.com/locate/resuscitation](http://www.elsevier.com/locate/resuscitation)EUROPEAN  
RESUSCITATION  
COUNCIL

## Ten Steps Toward Improving In-Hospital Cardiac Arrest Quality of Care and Outcomes ☆☆☆☆☆



**Paul S. Chan<sup>ag,\*,1</sup>, Robert Greif<sup>b,1</sup>, Theresa Anderson<sup>a</sup>, Huba Atiq<sup>d</sup>, Thomaz Bittencourt Couto<sup>e</sup>, Julie Considine<sup>f</sup>, Allan R. De Caen<sup>g</sup>, Therese Djärv<sup>h</sup>, Ann Doll<sup>i</sup>, Matthew J. Douma<sup>j</sup>, Dana P. Edelson<sup>k</sup>, Feng Xu<sup>l</sup>, Judith C. Finn<sup>m</sup>, Grace Firestone<sup>n</sup>, Saket Girotra<sup>o</sup>, Kasper G. Lauridsen<sup>p</sup>, Carrie Kah-Lai Leong<sup>q</sup>, Swee Han Lim<sup>q</sup>, Peter T. Morley<sup>r</sup>, Laurie J. Morrison<sup>s</sup>, Ari Moskowitz<sup>t</sup>, Ajit Mulasari Sankardas<sup>u</sup>, Mahmoud Tageldin Mustafa Mohamed<sup>v</sup>, Michelle Christy Myburgh<sup>w</sup>, Vinay M. Nadkarni<sup>x</sup>, Robert W. Neumar<sup>c</sup>, Jerry P. Nolan<sup>y</sup>, Justine Athieno Odakha<sup>z</sup>, Theresa M. Olasveengen<sup>aa</sup>, Judit Orosz<sup>ab</sup>, Gavin D. Perkins<sup>y</sup>, Jeanette K. Previdi<sup>ac</sup>, Christian Vaillancourt<sup>ad</sup>, William H. Montgomery<sup>ae</sup>, Comilla Sasson<sup>af,2</sup>, Brahmajee K. Nallamothu<sup>a,2</sup>, on behalf of the International Liaison Committee on Resuscitation**

**Keywords:** Adult, Cardiopulmonary resuscitation, Child, Humans, Incidence, Sudden cardiac arrest, Survival rate

Improving in-hospital cardiac arrest (IHCA) quality of care for adult and pediatric patients—not simply survival—requires a comprehensive set of programs and actions. Ideally, these should be embedded in a system of care that (1) plans and prepares for IHCA, (2) prevents IHCA when avoidable, (3) delivers high-quality, guideline-based resuscitation, and (4) continuously evaluates and improves itself within a culture of person-centered care.

IHCA is a high-risk event among hospitalized patients of all ages worldwide that is associated with significant morbidity and mortality.<sup>1</sup> Estimates of its incidence vary across industrialized countries with rates in adults between 1.2 and 10 per 1000 hospital admissions,<sup>2–5</sup> which translates to ≈300 000 IHCA events in the United States each year with a reported survival rate to hospital discharge of ≈25%.<sup>5</sup> Although experiences outside of higher income countries are limited, reported data suggest high incidence rates of IHCA in

low- to middle-income countries like Uganda and China.<sup>6,7</sup> These events are medical emergencies that require immediate treatment by teams of interdisciplinary health care professionals to optimize outcomes. Not surprisingly, there is a significant burden on hospitals to create and maintain resuscitation systems that are able to identify IHCA, activate an emergency response, and deliver high-quality resuscitation. Yet despite these pressures, there is consistent evidence that the quality of care around IHCA remains suboptimal and varies across hospitals and countries.<sup>8–10</sup>

Given the variation in IHCA quality of care and outcomes, the International Liaison Committee on Resuscitation (ILCOR) launched an initiative to provide strategic guidance delineating critical steps to improve IHCA care. The member councils comprising ILCOR are the American Heart Association, European Resuscitation Council, Heart and Stroke Foundation of Canada, Australian and New Zealand

☆ This manuscript was sent to Dennis T. Ko, Senior Guest Editor, for review by expert referees, editorial decision, and final disposition.

☆☆ This article has been co-published in *Circulation: Cardiovascular Quality and Outcomes*.

☆☆☆ This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

\* Corresponding author at: Paul S Chan, MD, MSc. Department of Internal Medicine. Saint Luke's Mid America Heart Institute, Kansas City, Missouri, United States.

E-mail address: [pchan@saint-lukes.org](mailto:pchan@saint-lukes.org) (P.S. Chan).

<sup>1</sup> P.S. Chan and R. Greif shared first authorship.

<sup>2</sup> C. Sasson and B.K. Nallamothu shared senior authorship.

<https://doi.org/10.1016/j.resuscitation.2023.109996>

0300-9572/© 2023 The Authors. Published by Elsevier B.V. on behalf of European Resuscitation Council.



**Fig. 1 – Ten steps toward improving in-hospital cardiac arrest (IHCA) quality of care and outcomes as a cycle of continuous improvement.**

Committee on Resuscitation, Resuscitation Councils of Asia, Indian Resuscitation Council Federation, and the collaborating organization, International Federation of Red Cross. The Ten Steps Toward Improving In-Hospital Cardiac Arrest Quality of Care and Outcomes in this document (Fig. 1; Table 1) is the result of this effort and builds upon prior work in the space of out-of-hospital cardiac arrest care by the Global Resuscitation Alliance.<sup>11</sup> These steps represent the consensus of a Writing Group of >30 interprofessional experts drawn from various branches of medicine, nursing, and allied health care professions.

For clarity and mutual understanding, we define IHCA as any infant, pediatric, or adult patient who sustains a cardiac arrest within a hospital setting recognizing that, for some hospitals, this may include surrounding geographic areas on a campus (eg, lobby, parking structures) and nonpatient visitors (we do not address neonatal resuscitation in this document given its distinctive aspects). We appreciate that this decision means there may be heterogeneity in how IHCA is defined by each hospital, as nonpatient visitors are unique from hospitalized patients, and they are likely to sustain a cardiac arrest for different reasons. However, for this document, the Writing Group felt that what is common across IHCA is established systems are needed to care for these critical events. IHCA necessitates immediate and appropriate recognition and intervention, often including immediate cardiopulmonary resuscitation (CPR), defibrillation, and other guideline-based interventions. We acknowledge that with this definition of IHCA that can occur in any setting within the hospital or its campus, IHCA will be distinct from out-of-hospital cardiac arrest in terms of etiology (eg, hypoxia, infection), response, and outcomes.<sup>12</sup>

## SELECTION OF WRITING AND ADVISORY GROUPS

Members of the Writing Group come from diverse regions across the globe, offering a unique blend of perspectives and experiences from both low- and high-resource settings. These individuals were selected by relying on nominations from ILCOR and other expert groups in resuscitation care from around the world. We tried to balance the membership across nationalities, professions, demographics, and resource settings. To support this Writing Group and ensure a diverse set of voices were heard, we also had an Advisory Board of >30 members who served to provide additional feedback throughout the process.

## METHODOLOGY OF SELECTING THE TEN STEPS

The process for selecting materials to include in this document entailed several rounds of brainstorming, discussion, and refinement to ensure the final steps truly reflect a global, comprehensive, and practical approach to improving IHCA care. Throughout the process, special consideration was placed on low-resource settings and the unique challenges they may face. Although this document reflects expert consensus guidance on best practices and published evidence from ILCOR statements,<sup>13,14</sup> it does not contain a systematic literature review or incorporate a formal grading of evidence.

Via a virtual platform, the Writing Group met numerous times over 5 months followed by a 2-day in-person meeting (May 28–29, 2023)

**Table 1 – Summary of ten steps toward improving in-hospital cardiac arrest quality of care and outcomes.**

<b>Plan and prepare</b>	<p><b>Step 1: build and support governance and infrastructure for a resuscitation program:</b> create a governance structure with administrative leadership to advance resuscitation excellence. Empower local operational champions and ensure a robust quality improvement program.</p> <p><b>Step 2: collect data to measure and improve resuscitation processes and outcomes:</b> use data to assess education, training, performance, and outcomes across the continuum of resuscitation care and to devise and evaluate the impact of quality improvement efforts to ensure their sustainability. Consider participating in resuscitation registries to enable benchmarking and contribute to knowledge transfer.</p> <p><b>Step 3: implement effective education and training for resuscitation:</b> create and maintain resuscitation education and training programs for resuscitation, including (1) standardized, in-person resuscitation courses; (2) low-dose, high-frequency in-situ training; (3) team-based training and simulations; and (4) debriefing to learn from resuscitation events.</p>
<b>Prevent</b>	<p><b>Step 4: establish patient and family goals of treatment early and reassess often:</b> proactively and collaboratively engage patients and their families in dialogue regarding their preferences for resuscitation in the context of their values, comorbidities, frailty, and family needs. Recognize that there are patients for whom resuscitation is unwanted or unlikely to result in an outcome that is acceptable to the patient. Ideally, goals of treatment conversations would start in the community and be updated on admission and after any change in clinical condition.</p> <p><b>Step 5: stop preventable IHCA:</b> have systems (eg, early warning systems) and teams (eg, medical/rapid response teams) in place to enable early prediction, identification, and response to deteriorating patients. Provide effective communication to ensure a clear plan and safe transition of care.</p>
<b>Perform</b>	<p><b>Step 6: develop and deploy an effective resuscitation response system:</b> make available a hospital-wide resuscitation response system that is easily and rapidly activated. Ensure a high-quality resuscitation team that includes preassigned, experienced, and interdisciplinary health care professionals.</p> <p><b>Step 7: deliver guideline-based resuscitation care:</b> focus on essential components of guideline-based skills for basic life support, advanced life support, and pediatric life support. These include prompt initiation of high-performance CPR and defibrillation and the use of appropriate and cause-directed interventions. Avoid nonrecommended interventions and early termination of resuscitation.</p> <p><b>Step 8: deliver guideline-based post-cardiac arrest care:</b> empower multidisciplinary teams to provide guideline-based post-cardiac arrest care, throughout and beyond hospitalization. Use multiple modalities to inform prognostication. Consider organ donation, when appropriate.</p>
<b>Principles and culture</b>	<p><b>Step 9: implement a person-centered culture of excellence in care:</b> ensure there is informed, culturally appropriate communication with patients and family to support them and to understand their goals, values, and preferences. For cardiac arrest survivors, development of postdischarge plans with targeted resources (eg, primary care, rehabilitation) is crucial for recovery and survivorship. For families of nonsurvivors, offer grief and bereavement resources.</p> <p><b>Step 10: ensure the well-being of health care professionals:</b> there should be a holistic focus on the psychological, physical, and spiritual well-being of health care professionals. Caring for the care team ensures that treatment of IHCA patients remains at a high quality.</p>

CPR indicates cardiopulmonary resuscitation; and IHCA, in-hospital cardiac arrest.

in Brisbane, Australia, that included an iterative group discussion format for voting, which resulted in the Ten Steps included in this document. We created several workgroups examining cardiac arrest phases of care (processes of care before, during, and after cardiac arrest), as well as key topic areas of focus (metrics and data, ethics, education, tools and devices, and teams and systems). Independent workgroups completed semistructured reports back to a Steering Committee and the entire Writing Group. Then during the 2-day in-person meeting, we had a series of facilitated discussions to vote on themes that arose from these discussions. We voted over 3 cycles and presented results from the voting (central tendencies, dispersion) to the Writing Group between cycles. At the conclusion of the meeting, we identified the Ten Steps to provide practical guidance for hospitals and health systems that are shared in this document. The Advisory Group was consulted 2 × throughout this process, and at the conclusion, to ensure the steps were relevant, practical, and applicable to a broad population of patients. The goal of this inclusive and consultative process was to create a document that respects diversity, embraces inclusivity, and speaks to the realities of a wide range of health care environments from low- to high-resource settings.

## IMPLEMENTATION OF THE STEPS

Our collective commitment to enhancing IHCA care has been the driving force behind this work, underscoring the essential role of collaboration in addressing complex health challenges. Our intent is not to impose a one-size-fits-all approach; rather, we recognize that local context, considering unique needs, geographic concerns, and available resources, will guide the prioritization and application of the Ten Steps. In other words, the steps are not meant to represent a natural order or a hierarchical sequence for their introduction into a system. Rather, each institution or health system should use these steps to identify the areas of the greatest need in determining which steps they would prioritize to achieve the highest impact on their IHCA care. The goal is to assist hospitals and health systems to establish, support, reinforce, evaluate, and improve their resuscitation systems for IHCA, with the aim of enhancing survival outcomes, ensuring quality care, and fostering a continuous cycle of improvement (Figure).

Our goal here is to provide practical guidance to improve IHCA quality of care and outcomes. To implement the Ten Steps in this document effectively, each institution should first conduct a thorough

needs assessment of its existing infrastructure, including its teams, protocols, and resources, as well as its outcomes related to IHCA. Based on this appraisal, the institution can then use the suggested steps to identify gaps, set priorities, and create an action plan tailored to its context and resources. Such approaches may involve creating a multidisciplinary team committed to the improvement process, ensuring that the communication lines are open and developing a shared understanding of the objectives.

We cannot underscore enough the critical role of champions, resources, and administrative support in implementing these steps—themes that came up frequently during our discussions. Achieving excellence in IHCA care is not simply about applying new techniques or protocols; it involves fostering a culture that values continuous improvement, patient-centered care, and interdisciplinary teamwork. The need for adequate resources, coupled with a strong commitment from the administration, is integral to nurturing such a culture and ensuring sustained improvement. Our guidance in the Ten Steps places a significant emphasis on affordability, availability, and sustainability of IHCA care. Addressing these aspects requires expanding our scope to cover essential areas such as training, education, and postresuscitation care (including beyond hospital discharge). This includes developing innovative, cost-effective solutions and strengthening the human resources through capacity building and continuous training, particularly in low-resource settings.

Many insights came after robust discussion among the Writing and Advisory Groups around challenges different hospitals and health care professionals face with IHCA. As a result, this document offers unique perspectives on IHCA care by going beyond the traditional focus on survival outcomes emphasized in guidelines. For instance, addressing emotional needs of families of both survivors and nonsurvivors of IHCA arose as a vital component of comprehensive care that is now embedded in this guidance. Moreover, opening conversations regarding organ donation with family members, when appropriate, arose from our discussions. Lastly, we underscore the necessity of supporting health care professionals involved in these emotionally charged, high-stake situations to ensure their physical and psychological safety. Without caring for health care professionals, the entire system cannot function at the highest level over the long term.

Finally, this document is not intended to be static; it is meant to be a living guide that should change and evolve with new evidence, experience, innovative practices, and feedback from its users. We suggest using this document and the associated tools and case examples that will be posted online at [www.ilcor.org/in-hospital-cardiac-arrest](http://www.ilcor.org/in-hospital-cardiac-arrest), to help others with implementation of the Ten Steps. This will allow these steps to remain responsive and relevant to the changing needs of health care institutions and the diversity of the populations they serve. Your contribution in the form of insights, experiences, suggestions, and case examples to this online site will help ensure the ongoing refinement and enhancement of these guidelines.

### **Step 1: Build and Support Governance and Infrastructure for a Resuscitation Program**

Create a governance structure with administrative leadership to advance resuscitation excellence. Empower local operational champions and ensure a robust quality improvement program.

#### **Summary**

Building a robust governance structure with administrative leadership committed to advancing resuscitation excellence is a key foundation

to improving survival from IHCA. Delivery of resuscitation care requires structures (eg, appropriate equipment, adequately trained health care professionals, prospective resuscitation database) and processes (eg, evidence-informed protocols, policies, and procedures). These integrate to form an effective system of resuscitation care, which leads to improved patient outcomes, held together by a framework of continuous quality improvement.<sup>15,16</sup>

#### **Strategies for Implementation**

Local operational champions leading resuscitation programs should be individuals who are invested in and have accountability and authority to oversee such programs, so that they can advocate for change and ensure the sustainability of changes and developments.<sup>17</sup> A recent study on hospital champions and IHCA survival found that hospitals with an active physician champion were 4× more likely to be in a higher survival quintile.<sup>18</sup> A review of champions in health care–related implementation consistently found that they are critical to implementation effectiveness.<sup>19</sup> Champions can engage administrative leadership to obtain resources needed for appropriate equipment (eg, defibrillators, resuscitation medications, airway devices, CPR compression quality-monitoring devices), advocate for participation in a resuscitation database, and ensure adequately trained health care professionals, to implement evidence-informed protocols, policies, and procedures.

Supportive infrastructure and policies will be needed for achieving high-quality care and outcomes in IHCA, to identify motivated and passionate champions, and to provide champions with resources to innovate and direct continuous quality improvement. Effective leaders and teams will draw from multiple disciplines and hospital departments (eg, medical, nursing, respiratory therapy, pharmacy) linked to best practices in IHCA education and training.<sup>20</sup> Champions who are present, visible, and who actively seek feedback and address concerns will be those who most effectively build trust within their teams and the larger resuscitation system.

### **Step 2: Collect Data to Measure and Improve Resuscitation Processes and Outcomes**

Use data to assess education, training, performance, and outcomes across the continuum of resuscitation care and to devise and evaluate the impact of quality improvement efforts to ensure their sustainability. Consider participating in resuscitation registries to enable benchmarking and contribute to knowledge transfer.

#### **Summary**

You cannot improve what you do not measure. Measurement is the cornerstone of performance management and quality improvement. At a basic level, this should include measuring cardiac arrest occurrence and survival outcomes. However, collection of data across the broader continuum of in-hospital resuscitation care enables assessment of performance and identification of areas for improvement. The specific nature of the data to be collected should be driven by the resources and needs of the hospital that can differ across institutions. For example, measurement of CPR quality could provide insights regarding CPR training needs<sup>21</sup>; measurement of time to defibrillation could identify the need for easier and faster access to defibrillators.<sup>22</sup>

The ILCOR Utstein reporting statement for IHCA represents one example of how hospitals and health systems could collect such data. This statement provides guidance for developing core and supplemental data elements across 6 domains: hospital, patients, pre-

event, cardiac arrest process, postresuscitation process, and outcomes.<sup>23</sup> To enable comparisons both within and between institutions, adherence to Utstein definitions for variables optimizes usefulness of the data collected.<sup>23</sup>

Over the last decade, there has been a proliferation of national and regional out-of-hospital cardiac arrest registries.<sup>24</sup> Although participation in a cardiac arrest registry can help catalyze quality improvement efforts through benchmarking, there are relatively few national or regional IHCA registries.<sup>25</sup> Exceptions include the American Heart Association Get With The Guidelines-Resuscitation registry,<sup>26</sup> the National Cardiac Arrest Audit in the United Kingdom,<sup>27</sup> and several smaller registries.<sup>1</sup> There is evidence from observational studies that Get With The Guidelines-Resuscitation participation duration is associated with a significant improvement in IHCA quality of care<sup>28</sup> and event survival.<sup>29,30</sup>

### *Strategies for Implementation*

The goal of IHCA data collection is to inform quality improvement initiatives. As such, it is important that there is buy-in from hospital leadership and resuscitation stakeholders, given that data collection requires resources and personnel (software, infrastructure, and dedicated personnel). The extent of data collection efforts (eg, number of data elements) can be tailored to institutional priorities and available resources. For hospitals with limited resources, it is reasonable to start with collecting core variables<sup>23</sup> initially and escalating data collection efforts over time.<sup>1</sup>

There are 2 important hospital data collection issues. First, identifying patients who are at risk for or experience an IHCA event is not straightforward, and institutions should develop a consistent process to label all IHCA cases (eg, based on the Utstein definition). Second, collecting IHCA data in real time is often challenging as it can interfere with resuscitation care delivery. Moreover, data collection hinges on accuracy of documentation by frontline professionals. Therefore, institutions should develop teams trained to collect IHCA data, as well as harmonizing hospital information systems for supplementing data elements for the registry. Furthermore, strategies to follow-up patients after hospital discharge should be considered to better understand long-term outcomes.

### **Step 3: Implement Effective Education and Training for Resuscitation**

Create and maintain resuscitation education and training programs for resuscitation, including (1) standardized, in-person resuscitation courses; (2) low-dose, high-frequency in-situ training; (3) team-based training and simulations; and (4) debriefing to learn from resuscitation events.

#### *Summary*

To improve patient outcomes, resuscitation education and training programs are essential for all hospital staff. Educational programs should be guideline based and tailored to meet employees' needs.<sup>31–34</sup> Knowledge may be acquired and retained using didactic sessions, blended learning, or online learning formats.<sup>21,35</sup> Staff training should emphasize team competencies, recognition of cardiac arrest, and incorporate the hands-on practice of psychomotor skills to deliver high quality with early defibrillation if indicated.

Education ideally should be spaced over time in a low-dose, high-frequency manner.<sup>36–39</sup> Programs may include skill stations to train psychomotor skills where only 2 to 5 minutes of hands-on training

every 1 to 4 months can improve skill acquisition and retention.<sup>40,41</sup> Contextual skills and teamwork competencies may be promoted using in-situ simulations tailored to the setting and the needs of health care professionals. In-situ simulations may be planned training sessions or unannounced to test the system and identify latent safety threats.<sup>42–44</sup> Debriefing after clinical cardiac arrests allows participants to reflect on resuscitation events, clinical performance, and goals for improvements and has been associated with improved patient outcomes.<sup>20</sup>

### *Strategies for Implementation*

An essential element to successful education and training for IHCA is strong and responsive leadership support with appropriate resource allocation (personnel, time, equipment, physical space, etc). Leadership should prioritize and promote resuscitation education and share data from clinical cardiac arrests.<sup>20</sup>

Developing an educational program starts with a need assessment to identify knowledge gaps and skill deficiencies among hospital staff. Next, designing a tailored program and piloting the intervention with a small cohort allows project leads to address issues before scaling to a larger audience. Include a train-the-trainer session to have highly skilled employees learn how to train larger groups. Integrate real-time feedback (ie, CPR feedback with devices or CPR coaches) and debriefing sessions into the training. Constructive feedback allows learners to reflect on self-performance and individualized areas for improvement.

Implement a robust evaluation and sustainability process to measure the effectiveness of the training program over time. Outcomes data on IHCA should be used to inform the process of continuous quality improvement.<sup>45</sup>

### **Step 4: Establish Patient and Family Goals of Treatment Early and Reassess Often**

Proactively and collaboratively engage patients and their families in dialogue regarding their preferences for resuscitation in the context of their values, comorbidities, frailty, and family needs. Recognize that there are patients for whom resuscitation is unwanted or unlikely to result in an outcome that is acceptable to the patient. Ideally, goals of treatment conversations would start in the community and be updated on admission and after any change in clinical condition.

#### *Summary*

Goals of treatment discussions play a crucial role in patient care, especially when addressing emergency treatment decisions like resuscitation.<sup>46,47</sup> Ideally, these discussions should occur before hospital admission and involve the patient and their primary care practitioner. These should consider realistic goals for treatment alongside patients' personal values and preferences and any advanced directives.<sup>48</sup> Upon admission to the hospital, it is essential to revisit and establish clear goals of treatment (eg, relevant options, limited time trials), as well as specifically addressing emergency treatments such as CPR, as some patients may not want to be resuscitated.<sup>49</sup>

During the hospital stay, it is necessary to reassess goals of treatment as the patient's illness or injury evolves, following a change in patient preferences or their risk of cardiac arrest, or when transferring to a different care setting.<sup>48,49</sup> This includes continuously evaluating the effectiveness of ongoing therapies and considering the option of withdrawing life-sustaining treatments. Overall, fostering

open and informed discussions about treatment goals between patients, their families, and health care professionals is vital to delivering patient-centered care and ensuring that medical decisions are in line with the patient's preferences and values.

### *Strategies for Implementation*

Structured communication tools improve the quality of communication, patient involvement, and documentation of patient wishes and treatment plans.<sup>50–52</sup> Staff need to be supported with training in the use of such tools, and recommendations should be recorded to ensure that these decisions are accessible and transferrable within and between care settings. If organ donation is appropriate, ensure trained staff handle the interface with the patient's family and the treatment team is made aware.<sup>53</sup>

Culturally sensitive materials should be developed and implemented to acknowledge that approaches to goals of treatment will vary across communities. Collaborative decision-making models that involve families, community leaders, and health care professionals can be used to ensure that cultural values and preferences are respected while honoring the individual's wishes (eg, family presence during resuscitation).

### **Step 5: Stop Preventable IHCA**

Have systems (eg, early warning systems) and teams (eg, medical/rapid response teams) in place to enable early prediction, identification, and response to deteriorating patients. Provide effective communication to ensure a clear plan and safe transition of care.

#### *Summary*

Preventing IHCA will reduce hospital morbidity and mortality. While not all IHCA are predictable or avoidable, having an approach to stopping preventable IHCA is a key step toward saving lives. Stopping preventable IHCA requires the establishment of afferent alert and efferent rapid response systems that are bolstered by robust implementation, governance, and continuous quality improvement. Reliable methods to communicate treatment plans and patient values or preferences among clinical teams are critical in stopping preventable IHCA and making appropriate resuscitation interventions (including not starting a resuscitation).

Published rates of preventable IHCA range from 9% to 78% in hospital wards and from 10% to 41% in intensive care units.<sup>54–59</sup> Opportunities for addressing preventable IHCA should be prioritized by local hospital cardiac arrest and quality improvement committees.<sup>56,60,61</sup> The ILCOR Consensus on Science With Treatment Recommendations suggests that hospitals consider the introduction of a rapid response system to support deteriorating patients and reduce the incidence of IHCA.<sup>52</sup> However, a recent Cochrane review reported that early warning scoring and rapid response systems provide little or no benefit.<sup>63</sup> Despite limited evidence that early warning scoring and rapid response systems improve outcomes, they are a global patient safety mainstay and recommended by several key organizations.<sup>15,64,65</sup>

### *Strategies for Implementation*

Implementation of rapid response systems is complex, and data on best practices are limited.

Evidence suggests that high-performing rapid response systems are staffed by a multidisciplinary team of dedicated personnel without competing responsibilities, who work collaboratively with bedside

teams, and foster a culture of safety that enables system activation without concern for reprisal.<sup>66</sup> Rapid response systems should work 24/7, which presents instances where shared decision-making might be difficult. In such circumstances, transfer of patients to a high level of care may provide time for involvement of other teams in decisions on resuscitation. Rapid response systems might also consider palliative and end-of-life care during a response.<sup>67–69</sup>

Because escalation of care to prevent IHCA is often driven by vital sign abnormalities or clinician concern, regular measurement and interpretation of vital signs is critical to IHCA prevention. Severity-of-illness scoring systems can be used for assessment and detection of deterioration, thus prompting a timely clinical response. There is increasing interest in integrating machine learning–based and artificial intelligence–powered systems to enable early detection of the deteriorating patient.<sup>70,71</sup>

### **Step 6: Develop and Deploy an Effective Resuscitation Response System**

Make available a hospital-wide resuscitation response system that is easily and rapidly activated. Ensure a high-quality resuscitation team that includes preassigned, experienced, and interdisciplinary health care professionals.

#### *Summary*

Even a minute of delay can significantly worsen a cardiac arrest victim's likelihood of survival.<sup>22,72,73</sup> Prompt detection of cardiac arrest and delivery of basic life support including high-quality CPR, defibrillation, and timely treatment by initial responders are critical. Immediate activation and response from a multidisciplinary resuscitation team that can initiate advanced life support is essential. High-performing systems ensure that resuscitation teams are supported by administrative leadership.

Over the last several decades, there has been substantial emphasis on training health care staff on resuscitation protocols.<sup>74</sup> Recent studies have highlighted strategies for implementing resuscitation teams at hospitals that consistently achieve high IHCA survival.<sup>9,17,20,66,75</sup> These attributes include dedicated or designated resuscitation teams that are preassigned, multidisciplinary team members with clear roles and responsibilities, and effective communication and leadership during IHCA. Moreover, a strong emphasis on training, mentoring, and empowerment of frontline health care professionals like nurses was also present in these studies.

### *Strategies for Implementation*

We propose that hospitals implement a resuscitation team that can be rapidly deployed to a patient in cardiac arrest. Emergency plans to bring equipment such as a defibrillator, emergency drugs, and airway to the bedside are vital. All initial responders should be trained to start CPR.

To address the complex and myriad needs of patients with IHCA, the resuscitation team should ideally include medical staff and nurses with other health professionals as appropriate. Resuscitation teams are more effective when members are preassigned and available to respond immediately.<sup>9,76</sup> At some institutions, hospital staff who respond to cardiac arrest and other hospital emergencies have limited-to-no other responsibilities.<sup>66</sup> While such a model may not be universally applicable, it is important that the team leader and key roles such as who will perform chest compressions, provide defibrillation, and manage the airway are delineated before, or early during a cardiac arrest, and matched to individual skills. Strong effective

leadership, closed-loop communication, and mutual respect are hallmarks of highly effective resuscitation teams. It is also vital that resuscitation teams are well supported with education and training efforts that go beyond traditional training on cardiac arrest algorithms.<sup>20</sup>

### **Step 7: Deliver Guideline-Based Resuscitation Care**

Focus on essential components of guideline-based skills for basic life support, advanced life support, and pediatric life support. These include prompt initiation of high-performance CPR and defibrillation and the use of appropriate and cause-directed interventions. Avoid nonrecommended interventions and early termination of resuscitation.

#### *Summary*

Global consensus has been achieved through ILCOR and the regional resuscitation councils on what constitutes best practice for the care of a patient with IHCA. Adherence to guidelines for IHCA is associated with improved rates of return of spontaneous circulation (ROSC), survival to hospital discharge, and long-term neurological outcome and quality of life.<sup>77,78</sup> Key components of resuscitation care include prompt initiation of high-performance CPR, defibrillation (when indicated), and the use of appropriate and cause-directed interventions to address the etiology of the IHCA. Efforts also should be made to avoid nonrecommended drugs, interventions, and early termination of resuscitation.<sup>13,79–81</sup>

#### *Strategies for Implementation*

As soon as an IHCA event is detected, start CPR immediately. Efforts should be made to closely monitor CPR quality by assessing the rate, depth, recoil, and compression fraction. This could be assisted by the use of capnography or a CPR feedback device. Rotate the individual performing chest compressions every 2 minutes or earlier, minimize interruptions to maintain quality of chest compressions (eg, during peri-shock pause, airway management, ventilations, and rhythm analysis), and optimize hand position during compression (lower half of sternum).

Initiate rhythm analysis and perform manual defibrillation (when indicated) as quickly as possible, ideally within 2 to 3 minutes of detecting cardiac arrest as each minute of delay is associated with lower survival.<sup>22</sup> Defibrillators should be user-friendly and chosen based on the probable users. Place defibrillators strategically in hospital to ensure quick access.

Monitor and optimize ventilation, considering factors like rate and volume, while avoiding hyperventilation. Whenever possible, bag-mask ventilation is ideally performed by 2 providers as 1-person bag-mask ventilation can be challenging. Avoid interruptions to chest compressions when inserting a supraglottic airway device or tracheal tube. An end-tidal carbon dioxide waveform capnography device should be used to confirm and monitor tracheal tube placement,<sup>82</sup> as classic clinical signs of tracheal intubation are not reliable.<sup>83</sup>

Inject intravenous epinephrine (adrenaline) as quickly as possible for IHCA with a nonshockable rhythm (and ideally within the first 5 minutes)<sup>73</sup> and after the second or third defibrillation for IHCA with a shockable rhythm.<sup>84</sup> Repeat epinephrine at 3- to 5-minute intervals. Consider the intraosseous route if unable to obtain intravenous access. Administer lidocaine or amiodarone for shock-refractory ventricular fibrillation. Do not give calcium or bicarbonate unless there

are specific indications (eg, hyperkalemia, tricyclic antidepressant overdose).

Avoid premature termination of resuscitation efforts.<sup>85,86</sup> In the absence of clear termination guidelines, data-informed standardized institutional decision algorithms should be developed collaboratively with consensus from all resuscitation stakeholders.

### **Step 8: Deliver Guideline-Based Postcardiac Arrest Care**

Empower multidisciplinary teams to provide guideline-based postcardiac arrest care, throughout and beyond hospitalization. Use multiple modalities to inform prognostication. Consider organ donation, when appropriate.

#### *Summary*

Post-cardiac arrest care begins as soon as ROSC is established and should be delivered by a multidisciplinary team to address components of pathophysiology, psychology, and stress syndromes. Team composition may be determined by the clinical context and available resources. Team members should have ready access to and a clear understanding of contemporary guidelines and their institution's protocols upon which to base care. Guideline-based care continues beyond the intensive care phase and into the ward/rehabilitation hospital/home setting and should engage, monitor, and support family and community as team members.<sup>87,88</sup> Palliative care and organ donation services should be involved as needed based upon individual patient needs and clinical context.

#### *Strategies for Implementation*

Accurately assessing and communicating the patient's prognosis at various intervals after ROSC helps families and caregivers prepare for potential outcomes and informs decisions about the relative benefits and risks of life-sustaining treatments. Population statistics based on patient demographic and arrest characteristics, especially when based on a hospital's own data, can provide families and caregivers a general idea of the patient's prognosis. However, the accuracy of prognostication at the individual level is limited in the early period after ROSC and typically is deferred (eg, for at least 72 hours after ROSC) under current guidelines. The accuracy of prognostication is also dependent on the modalities used, when they are measured, and potential confounding factors. It is recommended that hospitals use > 1 validated predictor of poor neurological outcome when making decisions about limiting life-sustaining therapy as no modality is 100% accurate.

Patients resuscitated from out-of-hospital cardiac arrest who meet circulatory or brain death criteria or other irrecoverable injury are potential multiple-organ donors.<sup>89–91</sup> While a knowledge gap exists with respect to organ recovery following IHCA,<sup>92</sup> there is the potential of appropriate referrals for organ donation for patients with circulatory or brain death after achieving ROSC,<sup>93</sup> recognizing there is variation in perceptions of death in relation to organ recovery,<sup>94</sup> as well as cultural and legal considerations.<sup>53</sup>

### **Step 9: Implement a Person-Centered Culture of Excellence in Care**

Ensure there is informed, culturally appropriate communication with patients and family to support them and to understand their goals, values, and preferences. For cardiac arrest survivors, development of postdischarge plans with targeted resources (eg, primary care, rehabilitation) is crucial for recovery and survivorship. For families of nonsurvivors, offer grief and bereavement resources.

### Summary

The involvement and influence of family members for hospitalized patients vary across different cultures and environments. Relatives might be present during a resuscitation and be advocates for patients during critical phases of care, and their involvement is integral to patients' well-being.<sup>95,96</sup> Family-centered care during IHCA embraces dedicated policies that enhance patient well-being, including consideration of the family context, illness-specific education for staff, and collaboration between patient, family, and health care professionals.<sup>97</sup>

Patients' wishes for treatment may change depending on their clinical condition and the context around them.<sup>98</sup> Thus, it is important to ensure that patients and families are aware of the current goals of treatment, to address hopes and fears proactively and to engage patients and family members repeatedly.<sup>99</sup>

Despite resuscitation efforts, patients sustaining an IHCA may not survive, so it is important to facilitate a good death, where end-of-life care is a crucial part of patient- and family-centered care. Meeting personal comfort needs, such as having a nonhospital blanket or food from home, creates peaceful memories during final moments<sup>100,101</sup> and may play a role in the emotional well-being of family members who often experience depression, complicated grief, and reduced quality of life.<sup>100</sup> Families of nonsurvivors of IHCA need support for grief and bereavement.<sup>102,103</sup>

Survivorship from IHCA comes with many emotions, and survivors may struggle to develop their own adaptive strategies to continue their lives. Therefore, survivors need tailored rehabilitation and recovery plans based on what is important to them.<sup>104</sup> Comprehensive discharge plans should encompass physical, cognitive, psychological, and social goals beyond those specifically related to IHCA.<sup>105</sup> Connection with other survivors through established social networks can facilitate peer-to-peer interactions with others who share a similar life experience.

### Strategies for Implementation

Health care professionals should plan regular meetings with patients and families where hopes and fears, as well as current goals of treatment, are on the agenda. A key part of this process will be to ensure that adequate social support services are provided to families. It would be ideal for providers to offer written information for anything that needs to be remembered or acted upon. For families of nonsurvivors, it is important to offer counseling support for grief when needed.

Overall, it would be ideal for hospitals to have a long-term discharge plan encompassing all aspects of survivors' lives including referral to appropriate physicians and support groups. This should include not only the physical sequelae of IHCA (physical and occupational therapy) but also interventions to address mental health (eg, posttraumatic stress disorder, depression, anxiety) and access to such care.

### Step 10: Ensure the Well-Being of Health Care Professionals

There should be a holistic focus on the psychological, physical, and spiritual well-being of health care professionals. Caring for the care team ensures that treatment of IHCA patients remains at a high quality.

### Summary

The emotional and physical demands of resuscitations, coupled with the potential for unfavorable outcomes, can result in significant psy-

chological burden on those involved. This may lead to stress, feelings of inadequacy and failure, burnout, and posttraumatic stress disorder that may persist long term.<sup>106–109</sup> These sequelae are a threat to health care professionals' well-being, staffing, quality of care, and stability of the health care system.<sup>110</sup>

Local champions and administrators should routinely assess the emotional state of their staff (and themselves) and offer appropriate psychological support, as needed (eg, therapists or psychiatrists, mentorship, and mindfulness meditation).<sup>111</sup> In addition, spiritual support should be considered and may take form of having a postresuscitation moment of silence, providing physical space for prayer and meditation, or involvement of chaplain staff.<sup>112</sup> These interventions should take diversity of beliefs, traditions, and cultures into account.

Coaching and mentorship for health care professionals are additional methods for ensuring provider well-being.<sup>113–115</sup> Peer mentorship can provide essential emotional support, assist with coping strategies, and facilitate psychological resilience amid acute traumatic events.<sup>116,117</sup> Sitting down to check in with a peer after a resuscitation is one way to acknowledge the experience of that individual.

Debriefings after IHCA may be used for quality improvement or psychological well-being.<sup>118,119</sup> Debriefings held immediately after the arrest (hot debriefings)<sup>120</sup> may be conducted to check on care providers' emotional reactions.<sup>121</sup> Debriefings can also be used to assess health care professionals' well-being and identify staff members in need of additional support. Processes should be in place to ensure any identified staff receive necessary services with guarantees of confidentiality.

### Strategies for Implementation

Initiatives that address the psychological stress and trauma of caring for IHCA patients require strong administrative and leadership support. Program development and sustainability requires dedicated resources for staff time and an infrastructure to guarantee confidentiality and psychological safety. Staff members should have input into the program at every level. Factors for successful implementation include rigorous program evaluation, a growth mindset culture, buy-in by staff and leadership, ensuring sufficient resource allocation, and prioritization of goals.<sup>122</sup>

## CONSIDERATIONS FOR LOW-RESOURCE SETTINGS

Providing resuscitation care in low-resource settings presents unique challenges, including access to training in life support skill courses, sufficient staffing, appropriate equipment and technology, funding for data collection and quality improvement, competing medical priorities to IHCA for hospitals to consider, and a lack of experienced leadership in this area. Nonetheless, core elements of many of the Ten Steps can be adapted to the local clinical context, bearing in mind cultural differences and resource limitations. We share some of these unique aspects and principles below.

One key principle was that starting with modest steps may require minimal investment of time and resources but have significant clinical impact (eg, identifying a local champion). Professional organizations and societies can be a resource for expertise and practical advice for initiating resuscitation programs in low-resource set-



tings to maximize their effectiveness. They can provide assistance with prioritization so that high-impact, low-cost processes are used to kick-start new programs.<sup>123</sup> For example, IHCA data collection and outcome measurements may be linked with ongoing quality improvement processes in countries or areas with limited resources. Institutional or single-center IHCA registries may be helpful in understanding local factors and unique challenges in such settings.

Another key principle is to consider varying cultural approaches and religious beliefs in different areas of the world, which affect IHCA, especially in low-resource settings. Patients and families expectations with regard to advanced directives, treatment escalation, or termination of care might be different from known evidence, which might put enormous pressure on the resuscitation teams. Cultural beliefs could limit the conduct of CPR by personnel of the opposite sex. Public awareness campaigns and health care education programs in hospital waiting areas might be a place to inform the broader public about resuscitation care. Indeed, some areas of the world may be unfamiliar with resuscitation leading to confusion (and even anger) when health care professionals initiate basic treatments like chest compressions. Collaborative decision-making models that involve families, community leaders, and health care professionals can be used to ensure that education for patients and families is available while cultural values and preferences are respected to honor the individual's wishes.

Finally, another principle is how resuscitation care training is implemented. For instance, adequate hospital staff training in low-resource settings might be focused first on a core group of trained responders and qualified personnel (eg, those in high-risk units like intensive care areas). This could take the form of low-dose, high-frequency training implemented in daily work routines and utilize local resources for deploying makeshift mannequins. Finally, e-learning and remote learning opportunities may provide a cost-effective way to obtain and retain knowledge for health care professionals in low-resource settings.

For more detailed discussion on this topic, we refer readers to a recent document on cardiac arrest in low-resource settings supported by ILCOR and several professional organizations.<sup>123</sup>

---

## SUMMARY

The Ten Steps Toward Improving In-Hospital Cardiac Arrest Quality of Care and Outcomes provides a framework for all stakeholders to analyze their strengths and weaknesses in their response to all patients who sustain IHCA, regardless of age. They highlight critical areas in resuscitation care and suggest approaches to addressing components of these areas that will have the largest potential impact on lowering IHCA incidence and improving survival. These steps were identified as part of a comprehensive consensus process with interdisciplinary experts from diverse backgrounds and around the world. Yet it is important to remember that IHCA remains an understudied medical condition with significant knowledge gaps, and future research will be needed to improve quality of resuscitation care and outcomes after IHCA even further. In conclusion, the Ten Steps documented here reflect current knowledge of best practices within resuscitation systems and serve as a foundation for improving IHCA quality of care and outcomes across different settings.

---

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

---

## Acknowledgments

The authors would like to acknowledge as collaborators the following members of the Improving In-Hospital Cardiac Arrest Advisory Group: Jason Acworth, Hanan Fahad Alharbi, Dianne L. Atkins, Bernd W. Böttiger, Andrea Cortegiani, Keith Couper, Vihara Dasanayake, Jimena del Castillo, Maya Dewan, Rakesh Garg, Jan-Thorsten Gräsner, Takanari Ikeyama, Mukul Kapoor, Hiroshi Kurosawa, Candace N. Mannarino, Abel Martínez-Mejias, Tasuku Matsuyama, Namala Mkopi, Margaret Nicholson, Tatsuya Norii, Shinichiro Ohshimo, Gene Ong, Tia T. Raymond, Joseph Rossano, Georg M. Schmölzer, Federico Semeraro, Lokesh Tiwari, and Carolyn Zelop. The authors would also like to thank Amber Hoover and Chalan King for their administrative support throughout this project.

---

## Sources of Funding

This work received grant support from the Laerdal Foundation.

---

## Disclosures

Dr Chan has received research funding from the American Heart Association (AHA) and the National Heart, Lung, and Blood Institute (NHLBI). Dr Djärv is the chair of Swedish Resuscitation Council and the International Liaison Committee on Resuscitation (ILCOR) First Aid Task Force. A. Doll is employed by the Resuscitation Academy. Dr Edelson has equity interest and employment in AgileMD. Dr Girotra receives funding from the National Institutes of Health (R01HL160734, R56HL158803, and R01HL166305). He also receives a stipend from AHA for editorial work. Dr Greif is chair of ILCOR Education, Implementation and Team Task Force and the European Resuscitation Council board director of guidelines. Dr Lauridsen serves as an ERC Science and Education Committee member and member of the ILCOR EIT Task Force. Dr Moskowitz receives funding from NHLBI (R33HL162980). M.C. Myburgh serves on the ILCOR Pediatric Task Force. Dr Neumar is the ILCOR cochair. J.P. Nolan is the editor in chief of Resuscitation and receives National Institute for Health and Care Research (NIHR) grant funding. Dr Olasveengen is a board member of the Laerdal Foundation. Dr Perkins receives grant funding from the NIHR Applied Research Collaboration West Midlands, Resuscitation Council United Kingdom, Laerdal Foundation, and the British Heart Foundation and serves on the boards for the European Resuscitation Council, Resuscitation Council UK, ILCOR, and University Hospitals Coventry and Warwickshire. He also serves as an editor for Resuscitation and Resuscitation Plus. J.K. Previdi serves as the AHA Emergency Cardiovascular Care Committee chair and is an AHA product consultant. Dr Vaillancourt is a member of the Heart and Stroke Foundation of Canada Resuscitation Advisory Committee and receives funding from The Ottawa Hospital Academic Medical Organization.

Dr Montgomery serves as the ILCOR consultant. Dr Sasson is employed by AHA. Dr Nallamothu has been a principal investigator or coinvestigator on research grants from the NIH, AHA, Janssen, and Apple, Inc. He also receives compensation as an editor in chief of *Circulation: Cardiovascular Quality and Outcomes*. Finally, he is a coinventor on the US Utility Patent Number US15/356012 (US20170148158A1) entitled "Automated Analysis of Vasculature in Coronary Angiograms" that uses software technology with signal processing and machine learning to automate the reading of coronary angiograms, held by the University of Michigan. The patent is licensed to AngioInsight, Inc, in which he holds ownership shares and receives consultancy fees. Dr Nadkarni receives research funding to his institution from the National Institutes of Health, Agency for Healthcare Research and Quality, US Department of Defense, American Heart Association, Zoll Medical, RQI Partners, and Laerdal Foundation; he also serves as the President of the Society of Critical Care Medicine (SCCM) 2023-2024. The views expressed in the manuscript are his, and not intended to represent the views of the SCCM. The other authors report no conflicts.

## Author details

<sup>a</sup>Department of Internal Medicine, University of Michigan Medical, Ann Arbor, United States <sup>b</sup>Department of Anesthesiology and Pain Medicine, University of Bern, Switzerland <sup>c</sup>Department of Emergency Medicine, University of Michigan, Ann Arbor, United States <sup>d</sup>Centre of Excellence for Trauma and Emergencies, Aga Khan University Hospital, Pakistan <sup>e</sup>Institution as Hospital Israelita Albert Einstein, São Paulo, Brazil <sup>f</sup>Deakin University, Geelong, Australia <sup>g</sup>Division of Pediatric Critical Care, Stollery Children's Hospital, Edmonton, Canada <sup>h</sup>Department of Medicine, Karolinska Institute, Stockholm, Sweden <sup>i</sup>Global Resuscitation Alliance, Seattle, WA, United States <sup>j</sup>Department of Critical Care Medicine, University of Alberta, Canada <sup>k</sup>Department of Medicine, University of Chicago Medicine, IL, United States <sup>l</sup>Department of Emergency Medicine, Qilu Hospital of Shandong University, China <sup>m</sup>School of Nursing, Curtin University, Perth, Australia <sup>n</sup>Department of Family Medicine, University of California Los Angeles Health, Santa Monica, United States <sup>o</sup>Department of Medicine, University of Texas Southwestern Medical Center, Dallas, United States <sup>p</sup>Department of Medicine, Randers Regional Hospital, Denmark <sup>q</sup>Department of Emergency Medicine, Singapore General Hospital, Singapore <sup>r</sup>Department of Intensive Care, The University of Melbourne, Australia <sup>s</sup>Division of Emergency Medicine, University of Toronto, Ontario, Canada <sup>t</sup>Department of Medicine, Montefiore Medical Center, The Bronx, NY, United States <sup>u</sup>Madras Medical Mission, Chennai, India <sup>v</sup>Burjeel Hospital, Abu Dhabi, United Arab Emirates <sup>w</sup>Department of Pediatric Emergency Medicine, University of the Free State, South Africa <sup>x</sup>Department of Anesthesiology and Critical Care, Childrens Hospital of Philadelphia, PA, United States <sup>y</sup>University of Warwick, Coventry, United Kingdom <sup>z</sup>Mbarara University of Science and Technology, Uganda <sup>aa</sup>Oslo University Hospital and Institute of Clinical Medicine, University of Oslo, Norway <sup>ab</sup>Department of Medicine, The Alfred, Melbourne, Australia <sup>ac</sup>American Heart Association, Dallas, TX, United States <sup>ad</sup>Department of Emergency Medicine, University of Ottawa, Canada <sup>ae</sup>Straub Clinic and Hospital, Honolulu, HI, United States <sup>af</sup>American Heart Association, Denver, CO, United States <sup>ag</sup>Mid-America Heart Institute, Kansas City, MO, United States

## REFERENCES

- Andersen LW, Holmberg MJ, Berg KM, Donnino MW, Granfeldt A. In-hospital cardiac arrest: a review. *J Am Med Assoc* 2019;321:1200–10. <https://doi.org/10.1001/jama.2019.1696>.
- Nolan JP, Soar J, Smith GB, Gwinnutt C, Parrott F, Power S, Harrison DA, Nixon E, Rowan K. National Cardiac Arrest Audit. Incidence and outcome of in-hospital cardiac arrest in the United Kingdom National Cardiac Arrest Audit. *Resuscitation* 2014;85:987–92. <https://doi.org/10.1016/j.resuscitation.2014.04.002>.
- Andersen LW, Holmberg MJ, Lofgren B, Kirkegaard H, Granfeldt A. Adult in-hospital cardiac arrest in Denmark. *Resuscitation* 2019;140:31–6. <https://doi.org/10.1016/j.resuscitation.2019.04.046>.
- Ohbe H, Tagami T, Uda K, Matsui H, Yasunaga H. Incidence and outcomes of in-hospital cardiac arrest in Japan 2011–2017: a nationwide inpatient database study. *J Intensive Care* 2022;10:10. <https://doi.org/10.1186/s40560-022-00601-y>.
- Holmberg MJ, Ross CE, Fitzmaurice GM, Chan PS, Duval-Arnould J, Grossestreuer AV, Yankama T, Donnino MW, Andersen LW. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Annual incidence of adult and pediatric in-hospital cardiac arrest in the United States. *Circ Cardiovasc Qual Outcomes* 2019;12. e005580. <https://doi.org/10.1161/CIRCOUTCOMES.119.005580>.
- Ocen D, Kalungi S, Ejoku J, Luggya T, Wabule A, Tumukunde J, Kizwera A. Prevalence, outcomes and factors associated with adult in hospital cardiac arrests in a low-income country tertiary hospital: a prospective observational study. *BMC Emerg Med* 2015;15:23. <https://doi.org/10.1186/s12873-015-0047-0>.
- Wang C, Zheng W, Zheng J, Shao F, Zhu Y, Li C, Ma Y, Tan H, Yan S, Han X, et al. A national effort to improve outcomes for in-hospital cardiac arrest in China: the Baseline Investigation of Cardiac Arrest (BASIC-IHCA). *Resusc Plus.* 2022;11. 100259. <https://doi.org/10.1016/j.resplu.2022.10025>.
- Chan PS, Nallamothu BK. Improving outcomes following in-hospital cardiac arrest: life after death. *J Am Med Assoc* 2012;307:1917–8. <https://doi.org/10.1001/jama.2012.3504>.
- Nallamothu BK, Guetterman TC, Harrod M, Kellenberg JE, Lehrich JL, Kronick SL, Krein SL, Iwashyna TJ, Saint S, Chan PS. How do resuscitation teams at top-performing hospitals for in-hospital cardiac arrest succeed? A qualitative study *Circulation* 2018;138:154–63. <https://doi.org/10.1161/CIRCULATIONAHA.118.033674>.
- Penketh J, Nolan JP. In-hospital cardiac arrest: the state of the art. *Crit Care* 2022;26:376. <https://doi.org/10.1186/s13054-022-04247-y>.
- Global Resuscitation Alliance. Accessed June 12, 2023. <https://www.globalresuscitationalliance.org/ten-programs/>.
- Hoybye M, Stankovic N, Holmberg M, Christensen HC, Granfeldt A, Andersen LW. In-hospital vs out-of-hospital cardiac arrest: patient characteristics and survival. *Resuscitation* 2021;158:157–65. <https://doi.org/10.1016/j.resuscitation.2020.11.016>.
- Nolan JP, Maconochie I, Soar J, Olasveengen TM, Greif R, Wyckoff MH, Singletary EM, Aickin R, Berg KM, Mancini ME, et al. Executive summary 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A1–A22. <https://doi.org/10.1016/j.resuscitation.2020.09.009>.
- Nolan JP, Maconochie I, Soar J, Olasveengen TM, Greif R, Wyckoff MH, Singletary EM, Aickin R, Berg KM, Mancini ME, et al. Executive summary: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation* 2020;142(16\_suppl\_1):S2–S27. <https://doi.org/10.1161/CIR.0000000000000890>.
- Semeraro F, Greif R, Bottiger BW, Burkart R, Cimpoesu D, Georgiou M, Yeung J, Lippert F, S Lockey A, Olasveengen TM, European Resuscitation Council guidelines, et al. . European

- Resuscitation Council guidelines 2021: systems saving lives. *Resuscitation* 2021;161:80–97. <https://doi.org/10.1016/j.resuscitation.2021.02.008>.
16. Kronick SL, Kurz MC, Lin S, Edelson DP, Berg RA, Billi JE, Cabanas JG, Cone DC, Diercks DB, Foster JJ, et al. Part 4: systems of care and continuous quality improvement: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2015;132(18 suppl 2):S397–413. <https://doi.org/10.1161/CIR.0000000000000258>.
  17. O'Halloran A, Shepard LN, Nadkarni VM. Building a culture of champions: the importance of leadership in resuscitation. *J Am Heart Assoc* 2021;10. e020390. <https://doi.org/10.1161/JAHA.121.020390>.
  18. Chan JL, Lehrich J, Nallamothu BK, Tang Y, Kennedy M, Trumppower B, Chan PS. American Heart Association's Get With the Guidelines®-Resuscitation Investigators. Association between hospital resuscitation champion and survival for in-hospital cardiac arrest. *J Am Heart Assoc* 2021;10. e017509. <https://doi.org/10.1161/JAHA.120.017509>.
  19. Miech EJ, Rattray NA, Flanagan ME, Damschroder L, Schmid AA, Damush TM. Inside help: an integrative review of champions in healthcare-related implementation. *SAGE Open Med*. 2018;6. 2050312118773261. <https://doi.org/10.1177/2050312118773261>.
  20. Anderson TM, Secrest K, Krein SL, Schildhouse R, Guetterman TC, Harrod M, Trumppower B, Kronick SL, Pribble J, Chan PS, et al. Best practices for education and training of resuscitation teams for in-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes* 2021;14. e008587. <https://doi.org/10.1161/CIRCOUTCOMES.121.008587>.
  21. Cheng A, Nadkarni VM, Mancini MB, Hunt EA, Sinz EH, Merchant RM, Donoghue A, Duff JP, Eppich W, Auerbach M, et al; American Heart Association Education Science Investigators; and on behalf of the American Heart Association Education Science and Programs Committee, Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation; Council on Cardiovascular and Stroke Nursing; and Council on Quality of Care and Outcomes Research. Resuscitation education science: educational strategies to improve outcomes from cardiac arrest: a scientific statement from the American Heart Association. *Circulation*. 2018;138:e82–e122. <https://doi.org/10.1161/CIR.0000000000000583>.
  22. Chan PS, Krumholz HM, Nichol G, Nallamothu BK. American Heart Association National Registry of Cardiopulmonary Resuscitation Investigators. Delayed time to defibrillation after in-hospital cardiac arrest. *N Engl J Med* 2008;358:9–17. <https://doi.org/10.1056/NEJMoa0706467>.
  23. Nolan JP, Berg RA, Andersen LW, Bhanji F, Chan PS, Donnino MW, Lim SH, Ma MH, Nadkarni VM, Starks MA, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the utstein resuscitation registry template for in-hospital cardiac arrest: a consensus report from a task force of ILCOR. *Circulation* 2019;140:e746–57. <https://doi.org/10.1161/CIR.0000000000000710>.
  24. Nishiyama C, Kiguchi T, Okubo M, Alihodžić H, Al-Araji R, Baldi E, Beganton F, Booth S, Bray J, Christensen E, et al. Three-year trends in out-of-hospital cardiac arrest across the world: second report from the International Liaison Committee on Resuscitation (ILCOR). *Resuscitation* 2023;186. 109757. <https://doi.org/10.1016/j.resuscitation.2023.109757>.
  25. Chan PS, Kennedy KF, Girotra S. American Heart Association's Get With The Guidelines®-Resuscitation Investigators. Updating the model for risk-standardizing survival for in-hospital cardiac arrest to facilitate hospital comparisons. *Resuscitation* 2023;183. 109686. <https://doi.org/10.1016/j.resuscitation.2022.109686>.
  26. American Heart Association. Get With The Guidelines® Resuscitation. Accessed June 12, 2023. <https://www.heart.org/en/professional/quality-improvement/get-with-the-guidelines/get-with-the-guidelines-resuscitation>.
  27. National Cardiac Arrest Audit (NCAA). Accessed June 16, 2023. <https://www.icnarc.org/Our-Audit/Audits/Ncaa/About>.
  28. Starks MA, Dai D, Nichol G, Al-Khatib SM, Chan P, Bradley SM, Peterson ED. American Heart Association's Get With the Guidelines-Resuscitation Investigators. The Association of Duration of participation in get with the guidelines-resuscitation with quality of care for in-hospital cardiac arrest. *Am Heart J* 2018;204:156–62. <https://doi.org/10.1016/j.ahj.2018.04.018>.
  29. Bradley SM, Huszti E, Warren SA, Merchant RM, Sayre MR, Nichol G. Duration of hospital participation in Get With the Guidelines-Resuscitation and survival of in-hospital cardiac arrest. *Resuscitation* 2012;83:1349–57. <https://doi.org/10.1016/j.resuscitation.2012.03.014>.
  30. Girotra S, Nallamothu BK, Spertus JA, Li Y, Krumholz HM, Chan PS. American Heart Association Get with the Guidelines-Resuscitation Investigators. Trends in survival after in-hospital cardiac arrest. *N Engl J Med* 2012;367:1912–20. <https://doi.org/10.1056/NEJMoa1109148>.
  31. Lauridsen KG, Løfgren B, Brogaard L, Paltved C, Hvidman L, Krogh K. Cardiopulmonary resuscitation training for healthcare professionals: a scoping review. *Simul Healthc* 2021;17:170–82.
  32. Patocka C, Lockey A, Lauridsen KG, Greif R. Impact of accredited advanced life support course participation on in-hospital cardiac arrest patient outcomes: a systematic review. *Resusc Plus*. 2023;14. 100389. <https://doi.org/10.1016/j.resplu.2023.100389>.
  33. Elgohary M, Palazzo FS, Breckwoldt J, Cheng A, Pellegrino J, Schnaubelt S, Greif R, Lockey A. Blended learning for accredited life support courses - a systematic review. *Resusc Plus*. 2022;10. 100240. <https://doi.org/10.1016/j.resplu.2022.100240>.
  34. Lockey A, Patocka C, Lauridsen K, Finn J, Greif T; on behalf of the International Liaison Committee on Resuscitation Education, Implementation and Teams Task Force. Are cardiac arrest patient outcomes improved as a result of a member of the resuscitation team attending an accredited advanced life support course? Consensus on science with treatment recommendations. International Liaison Committee on Resuscitation (ILCOR) Education, Implementation and Teams Task Force; 2022. Accessed August 13, 2023. <https://costr.ilcor.org/document/>.
  35. Ali DM, Hisam B, Shaikat N, Baig N, Ong MEH, Epstein JL, Goralnick E, Kivela PD, McNally B, Razzak J. Cardiopulmonary resuscitation (CPR) training strategies in the times of COVID-19: a systematic literature review comparing different training methodologies. *Scand J Trauma Resusc Emerg Med* 2021;29:53. <https://doi.org/10.1186/s13049-021-00869-3>.
  36. Yeung J, Djarv T, Hsieh MJ, Sawyer T, Lockey A, Finn J, Greif Education R. Implementation and Team Task Force and Neonatal Life Support Task Force of the International Liaison Committee on Resuscitation (ILCOR). Spaced learning versus massed learning in resuscitation - a systematic review. *Resuscitation* 2020;156:61–71. <https://doi.org/10.1016/j.resuscitation.2020.08.132>.
  37. Niles D, Sutton RM, Donoghue A, Kalsi MS, Roberts K, Boyle L, Nishisaki A, Arbogast KB, Helfaer M, Nadkarni V. "Rolling refreshers": a novel approach to maintain CPR psychomotor skill competence. *Resuscitation* 2009;80:909–12. <https://doi.org/10.1016/j.resuscitation.2009.04.021>.
  38. Sutton RM, Niles D, Meaney PA, Aplenc R, French B, Abella BS, Lengetti EL, Berg RA, Helfaer MA, Nadkarni V. Low-dose, high-frequency CPR training improves skill retention of in-hospital pediatric providers. *Pediatrics* 2011;128:e145–51. <https://doi.org/10.1542/peds.2010-2105>.
  39. O'Leary A, Butler P, Fine JR. Dedicated chest compressor team: a quality improvement initiative to improve chest compression performance at in-hospital cardiac arrest events through quarterly training. *Resusc Plus*. 2023;13. 100361. <https://doi.org/10.1016/j.resplu.2023.100361>.
  40. Anderson R, Sebaldt A, Lin Y, Cheng A. Optimal training frequency for acquisition and retention of high-quality CPR skills: a

- randomized trial. *Resuscitation* 2019;135:153–61. <https://doi.org/10.1016/j.resuscitation.2018.10.033>.
41. Sullivan NJ, Duval-Arnould J, Twilley M, Smith SP, Aksamit D, Boone-Guercio P, Jeffries PR, Hunt EA. Simulation exercise to improve retention of cardiopulmonary resuscitation priorities for in-hospital cardiac arrests: a randomized controlled trial. *Resuscitation* 2015;86:6–13. <https://doi.org/10.1016/j.resuscitation.2014.10.021>.
  42. Stærk M, Lauridsen KG, Støtt CT, Riis DN, Løfgren B, Krogh K. In-hospital cardiac arrest - the crucial first 5 min: a simulation study. *Adv Simulation (Lond)*. 2022;7:29. <https://doi.org/10.1186/s41077-022-00225-0>.
  43. Stærk M, Lauridsen KG, Johnsen J, Løfgren B, Krogh K. In-situ simulations to detect patient safety threats during in-hospital cardiac arrest. *Resusc Plus*. 2023;14. 100410. <https://doi.org/10.1016/j.resplu.2023.100410>.
  44. Rubio-Gurung S, Putet G, Touzet S, Gauthier-Moulinier H, Jordan I, Beissel A, Labaune JM, Blanc S, Amamra N, Balandras C, et al. In situ simulation training for neonatal resuscitation: an RCT. *Pediatrics* 2014;134:e790–7. <https://doi.org/10.1542/peds.2013-3988>.
  45. Ruangsomboon O, Surabenjawongse U, Jantataeme P, Chawaruechai T, Wangtawesap K, Chakorn T. Association between cardiopulmonary resuscitation audit results with in-situ simulation and in-hospital cardiac arrest outcomes and key performance indicators. *BMC Cardiovasc Disord* 2023;23:299. <https://doi.org/10.1186/s12872-023-03320-w>.
  46. Mentzelopoulos SD, Couper K, Voorde PV, Druwe P, Blom M, Perkins GD, Lulic I, Djakow J, Raffay V, Lilja G, et al. European Resuscitation Council guidelines 2021: ethics of resuscitation and end of life decisions. *Resuscitation* 2021;161:408–32. <https://doi.org/10.1016/j.resuscitation.2021.02.017>.
  47. Pitcher D, Fritz Z, Wang M, Spiller JA. Emergency care and resuscitation plans. *BMJ* 2017;356:j876. <https://doi.org/10.1136/bmj.j876>.
  48. Perkins GD, Hawkes CA, Eli K, Griffin J, Jacques C, Huxley CJ, Couper K, Ochieng C, Fuld J, Fritz Z, et al. Recommended summary plan for emergency care and treatment: ReSPECT a mixed-methods study. *Health Social Care Delivery Res* 2022;10:1–140. <https://doi.org/10.3310/lfpe3627>.
  49. Perkins GD, Griffiths F, Slowther AM, George R, Fritz Z, Satherley P, Williams B, Waugh N, Cooke MW, Chambers S, et al. Do-not-attempt-cardiopulmonary-resuscitation decisions: an evidence synthesis. *Health Social Care Delivery Res* 2016;4:1–154. <https://doi.org/10.3310/hsdr04110>.
  50. Field RA, Fritz Z, Baker A, Grove A, Perkins GD. Systematic review of interventions to improve appropriate use and outcomes associated with do-not-attempt-cardiopulmonary-resuscitation decisions. *Resuscitation* 2014;85:1418–31. <https://doi.org/10.1016/j.resuscitation.2014.08.024>.
  51. Curtis JR, Lee RY, Brumback LC, Kross EK, Downey L, Torrence J, LeDuc N, Mallon Andrews K, Im J, Heywood J, et al. Intervention to promote communication about goals of care for hospitalized patients with serious illness: a randomized clinical trial. *J Am Med Assoc* 2023;329:2028–37. <https://doi.org/10.1001/jama.2023.8812>.
  52. Becker C, Lecheler L, Hochstrasser S, Metzger KA, Widmer M, Thommen EB, Nienhaus K, Ewald H, Meier CA, Rueter F, et al. Association of communication interventions to discuss code status with patient decisions for do-not-resuscitate orders: a systematic review and meta-analysis. *JAMA Netw Open* 2019;2. e195033. <https://doi.org/10.1001/jamanetworkopen.2019.5033>.
  53. Morrison LJ, Sandroni C, Grunau B, Parr M, Macneil F, Perkins GD, Aibiki M, Censullo E, Lin S, Neumar RW, et al. International Liaison Committee on Resuscitation. Organ donation after out-of-hospital cardiac arrest: a scientific statement from the International Liaison Committee on Resuscitation. *Circulation* 2023;148:e120–46. <https://doi.org/10.1161/CIR.0000000000001125>.
  54. Moskowitz A, Berg KM, Cocchi MN, Chase M, Yang JX, Sarge J, Grossestreuer AV, Sarge T, O' Donoghue S, Donnino MW. Cardiac arrest in the intensive care unit: an assessment of preventability. *Resuscitation* 2019;145:15–20. <https://doi.org/10.1016/j.resuscitation.2019.09.003>.
  55. Galhotra S, DeVita MA, Simmons RL, Dew MA. Members of the Medical Emergency Response Improvement Team (MERIT) Committee. Mature rapid response system and potentially avoidable cardiopulmonary arrests in hospital. *Qual Saf Health Care* 2007;16:260–5. <https://doi.org/10.1136/qshc.2007.022210>.
  56. Hodgetts TJ, Kenward G, Vlackonikolis I, Payne S, Castle N, Crouch R, Ineson N, Shaikh L. Incidence, location and reasons for avoidable in-hospital cardiac arrest in a district general hospital. *Resuscitation* 2002;54:115–23. [https://doi.org/10.1016/s0300-9572\(02\)00098-9](https://doi.org/10.1016/s0300-9572(02)00098-9).
  57. Bingham G, Bilgrami I, Sandford M, Larwill S, Orosz J, Luckhoff C, Kambourakis T. Avoiding adult in-hospital cardiac arrest: a retrospective cohort study to determine preventability. *Aust Crit Care* 2018;31:219–25. <https://doi.org/10.1016/j.aucc.2017.05.002>.
  58. Fennessy G, Hilton A, Radford S, Bellomo R, Jones D. The epidemiology of in-hospital cardiac arrests in Australia and New Zealand. *Intern Med J* 2016;46:1172–81. <https://doi.org/10.1111/imj.13039>.
  59. ANZ-CODE Investigators. Antecedents to and outcomes for in-hospital cardiac arrests in Australian hospitals with mature medical emergency teams: a multicentre prospective observational study. *Aust Crit Care*. 2023;S1036-7314(23)00026-7. <https://doi.org/10.1016/j.aucc.2023.01.011>.
  60. Moskowitz A, Berg KM, Cocchi MN, Grossestreuer AV, Issa M, Balaji L, Chase M, Yang JX, Sarge J, O'Donoghue S, et al. A trigger and response system for preventing cardiac arrest in the ICU. *Crit Care Explor* 2021;3:e0557. <https://doi.org/10.1097/CCE.0000000000000557>.
  61. Lyons PG, Edelson DP, Carey KA, Twu NM, Chan PS, Peberdy MA, Praestgaard A, Churpek MM. American Heart Association's Get With the Guidelines – Resuscitation Investigators. Characteristics of rapid response calls in the United States: an analysis of the first 402,023 adult cases from the Get With the Guidelines Resuscitation-Medical Emergency Team Registry. *Crit Care Med* 2019;47:1283–9. <https://doi.org/10.1097/CCM.0000000000003912>.
  62. Greif R, Bhanji F, Bigham BL, Bray J, Breckwoldt J, Cheng A, Duff JP, Gilfoyle E, Hsieh MJ, Iwami T, et al. Education, implementation, and teams: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A188–239. <https://doi.org/10.1016/j.resuscitation.2020.09.014>.
  63. McGaughey J, Fergusson DA, Van Bogaert P, Rose L. Early warning systems and rapid response systems for the prevention of patient deterioration on acute adult hospital wards. *Cochrane Database Syst Rev* 2021;2021:Cd005529. <https://doi.org/10.1002/14651858.CD005529.pub3>.
  64. Australian Commission on Safety and Quality in Health Care. Essential Elements for Recognising and Responding to Acute Physiological Deterioration. 3rd ed. 2021. ISBN: 978-1-925665-95-6.
  65. Berg KM, Cheng A, Panchal AR, Topjian AA, Aziz K, Bhanji F, Bigham BL, Hirsch KG, Hoover AV, Kurz MC, et al. Adult Basic and Advanced Life Support, Pediatric Basic and Advanced Life Support, Neonatal Life Support, and Resuscitation Education Science Writing Groups. Part 7: systems of care: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2020;142(16\_suppl\_2):S580–604. <https://doi.org/10.1161/CIR.0000000000000899>.
  66. Dukes K, Bunch JL, Chan PS, Guetterman TC, Lehrich JL, Trumpower B, Harrod M, Krein SL, Kellenberg JE, Reisinger HS, et al. Assessment of rapid response teams at top-performing hospitals for in-hospital cardiac arrest. *JAMA Intern Med*

- 2019;179:1398–405. <https://doi.org/10.1001/jamaintermmed.2019.2420>.
67. Downar J, Barua R, Rodin D, Lejniaks B, Gudimella R, McCredie V, Hayes C, Steel A. Changes in end of life care 5 years after the introduction of a rapid response team: a multicentre retrospective study. *Resuscitation* 2013;84:1339–44. <https://doi.org/10.1016/j.resuscitation.2013.03.003>.
  68. Downar J, Rodin D, Barua R, Lejniaks B, Gudimella R, McCredie V, Hayes C, Steel A. Rapid response teams, do not resuscitate orders, and potential opportunities to improve end-of-life care: a multicentre retrospective study. *J Crit Care* 2013;28:498–503. <https://doi.org/10.1016/j.jcrc.2012.10.002>.
  69. Jones D, Moran J, Winters B, Welch J. The rapid response system and end-of-life care. *Curr Opin Crit Care* 2013;19:616–23. <https://doi.org/10.1097/MCC.0b013e3283636be2>.
  70. Escobar GJ, Liu VX, Schuler A, Lawson B, Greene JD, Kipnis P. Automated identification of adults at risk for in-hospital clinical deterioration. *N Engl J Med* 2020;383:1951–60. <https://doi.org/10.1056/NEJMsa2001090>.
  71. Winslow CJ, Edelson DP, Churpek MM, Taneja M, Shah NS, Datta A, Wang CH, Ravichandran U, McNulty P, Kharasch M, et al. The impact of a machine learning early warning score on hospital mortality: a multicenter clinical intervention trial. *Crit Care Med* 2022;50:1339–47. <https://doi.org/10.1097/CCM.0000000000005492>.
  72. Bircher NG, Chan PS, Xu Y. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Delays in cardiopulmonary resuscitation, defibrillation, and epinephrine administration all decrease survival in in-hospital cardiac arrest. *Anesthesiology* 2019;130:414–22. <https://doi.org/10.1097/ALN.0000000000002563>.
  73. Donnino MW, Saliccioli JD, Howell MD, Cocchi MN, Giberson B, Berg K, Gautam S, Callaway C. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Time to administration of epinephrine and outcome after in-hospital cardiac arrest with non-shockable rhythms: retrospective analysis of large in-hospital data registry. *BMJ* 2014;348:g3028. <https://doi.org/10.1136/bmj.g3028>.
  74. Carveth SW, Burnap TK, Bechtel J, McIntyre K, Donegan J, Buchman RJ, Reese HE. Training in advanced cardiac life support. *J Am Med Assoc* 1976;235:2311–5.
  75. Guetterman TC, Kellenberg JE, Krein SL, Harrod M, Lehrich JL, Iwashyna TJ, Kronick SL, Girotra S, Chan PS, Nallamothu BK. Nursing roles for in-hospital cardiac arrest response: higher versus lower performing hospitals. *BMJ Qual Saf* 2019;28:916–24. <https://doi.org/10.1136/bmjqs-2019-009487>.
  76. Lauridsen KG, Krogh K, Muller SD, Schmidt AS, Nadkarni VM, Berg RA, Bach L, Dodt KK, Maack TC, Moller DS, et al. Barriers and facilitators for in-hospital resuscitation: a prospective clinical study. *Resuscitation* 2021;164:70–8. <https://doi.org/10.1016/j.resuscitation.2021.05.007>.
  77. Crowley CP, Saliccioli JD, Kim EY. The association between ACLS guideline deviations and outcomes from in-hospital cardiac arrest. *Resuscitation* 2020;153:65–70. <https://doi.org/10.1016/j.resuscitation.2020.05.042>.
  78. Khara R, Tang Y, Link MS, Krumholz HM, Girotra S, Chan PS. Association between hospital recognition for resuscitation guideline adherence and rates of survival for in-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes* 2019;12. e005429. <https://doi.org/10.1161/CIRCOUTCOMES.118.005429>.
  79. Soar J, Maconochie I, Wyckoff MH, Olasveengen TM, Singletary EM, Greif R, Aickin R, Bhanji F, Donnino MW, Mancini ME, et al. 2019 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2019;145:95–150. <https://doi.org/10.1016/j.resuscitation.2019.10.016>.
  80. Wyckoff MH, Singletary EM, Soar J, Olasveengen TM, Greif R, Liley HG, Zideman D, Bhanji F, Andersen LW, Avis SR, et al. COVID-19 Working Group. 2021 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the Basic Life Support; Advanced Life Support; Neonatal Life Support; Education, Implementation, and Teams; First Aid Task Forces; and the COVID-19 Working Group. *Resuscitation* 2021;169:229–311. <https://doi.org/10.1016/j.resuscitation.2021.10.040>.
  81. Wyckoff MH, Greif R, Morley PT, Ng KC, Olasveengen TM, Singletary EM, Soar J, Cheng A, Drennan IR, Liley HG, et al; Collaborators. 2022 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the Basic Life Support; Advanced Life Support; Pediatric Life Support; Neonatal Life Support; Education, Implementation, and Teams; and First Aid Task Forces. *Resuscitation*. 2022;181:208–288. <https://doi.org/10.1016/j.resuscitation.2022.10.005>.
  82. Chrimes N, Higgs A, Hagberg CA, Baker PA, Cooper RM, Greif R, Kovacs G, Law JA, Marshall SD, Myatra SN, et al. Preventing unrecognised oesophageal intubation: a consensus guideline from the Project for Universal Management of Airways and International Airway Societies. *Anaesthesia* 2022;77:1395–415. <https://doi.org/10.1111/anae.15817>.
  83. Hansel J, Law JA, Chrimes N, Higgs A, Cook TM. Clinical tests for confirming tracheal intubation or excluding oesophageal intubation: a diagnostic test accuracy systematic review and meta-analysis. *Anaesthesia* 2023;78:1020–30. <https://doi.org/10.1111/anae.16059>.
  84. Evans E, Swanson MB, Mohr N, Boulos N, Vaughan-Sarrazin M, Chan PS, Girotra S. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Epinephrine before defibrillation in patients with shockable in-hospital cardiac arrest: propensity matched analysis. *BMJ* 2021;375:e066534. <https://doi.org/10.1136/bmj-2021-066534>.
  85. Goldberger ZD, Chan PS, Berg RA, Kronick SL, Cooke CR, Lu M, Banerjee M, Hayward RA, Krumholz HM, Nallamothu BK. American Heart Association Get With The Guidelines—Resuscitation (formerly National Registry of Cardiopulmonary Resuscitation) Investigators. Duration of resuscitation efforts and survival after in-hospital cardiac arrest: an observational study. *Lancet* 2012;380:1473–81. [https://doi.org/10.1016/S0140-6736\(12\)60862-9](https://doi.org/10.1016/S0140-6736(12)60862-9).
  86. Lauridsen KG, Djarv T, Breckwoldt J, Tjissen JA, Couper K, Greif Education R. Implementation and Team Task Force of the International Liaison Committee on Resuscitation (ILCOR). Pre-arrest prediction of survival following in-hospital cardiac arrest: a systematic review of diagnostic test accuracy studies. *Resuscitation* 2022;179:141–51. <https://doi.org/10.1016/j.resuscitation.2022.07.041>.
  87. Haywood K, Whitehead L, Nadkarni VM, Achana F, Beesems S, Böttiger BW, Brooks A, Castrén M, Ong ME, Hazinski MF, et al; COSCA Collaborators. COSCA (Core Outcome Set for Cardiac Arrest) in adults: an advisory statement from the International Liaison Committee on Resuscitation. *Circulation*. 2018;137:e783–e801. <https://doi.org/10.1161/CIR.0000000000000562>.
  88. Topjian AA, Scholefield BR, Pinto NP, Fink EL, Buysse CMP, Haywood K, Maconochie I, Nadkarni VM, de Caen A, Escalante-Kanashiro R, et al. P-COSCA (Pediatric Core Outcome Set for Cardiac Arrest) in children: an advisory statement from the International Liaison Committee on Resuscitation. *Circulation* 2020;142:e246–61. <https://doi.org/10.1161/CIR.0000000000000911>.
  89. Elmer J, Molyneaux BJ, Shutterly K, Stuart SA, Callaway CW, Darby JM, Weisgerber AR. Organ donation after resuscitation from cardiac arrest. *Resuscitation* 2019;145:63–9. <https://doi.org/10.1016/j.resuscitation.2019.10.013>.
  90. Ho AFW, Tan TXZ, Latiff E, Shahidah N, Ng YY, Leong BS, Lim SL, Pek PP, Gan HN, Mao DR, et al. Assessing unrealised potential for

- organ donation after out-of-hospital cardiac arrest. *Scand J Trauma Resusc Emerg Med* 2021;29:105. <https://doi.org/10.1186/s13049-021-00924-z>.
91. Renaudier M, Binois Y, Dumas F, Lamhaut L, Beganton F, Jost D, Charpentier J, Lesieur O, Marijon E, Jouven X, et al. Paris Sudden Death Expertise Center group. Organ donation after out-of-hospital cardiac arrest: a population-based study of data from the Paris Sudden Death Expertise Center. *Ann Intensive Care* 2022;12:48. <https://doi.org/10.1186/s13613-022-01023-7>.
  92. Sandroni C, D'Arrigo S, Callaway CW, Cariou A, Dragancea I, Taccone FS, Antonelli M. The rate of brain death and organ donation in patients resuscitated from cardiac arrest: a systematic review and meta-analysis. *Intensive Care Med* 2016;42:1661–71. <https://doi.org/10.1007/s00134-016-4549-3>.
  93. Witten L, Gardner R, Holmberg MJ, Wiberg S, Moskowitz A, Mehta S, Grossestreuer AV, Yankama T, Donnino MW, Berg KM. Reasons for death in patients successfully resuscitated from out-of-hospital and in-hospital cardiac arrest. *Resuscitation* 2019;136:93–9. <https://doi.org/10.1016/j.resuscitation.2019.01.031>.
  94. Skowronski G, Ramnani A, Walton-Sonda D, Forlini C, O'Leary MJ, O'Reilly L, Sheahan L, Stewart C, Kerridge I. A scoping review of the perceptions of death in the context of organ donation and transplantation. *BMC Med Ethics* 2021;22:167. <https://doi.org/10.1186/s12910-021-00734-z>.
  95. Othman EH, Al Omar S, Abualruz H, Safadi R. An Islamic cultural perspective of family presence during resuscitation. *Resuscitation* 2020;151:75–6. <https://doi.org/10.1016/j.resuscitation.2020.04.007>.
  96. Afzali Rubin M, Svensson TL, Herling SF, Jabre P, Moller AM. Family presence during resuscitation. *Cochrane Database Syst Rev* 2023;5:CD013619. <https://doi.org/10.1002/14651858.CD013619.pub2>.
  97. Kokorelias KM, Gignac MAM, Naglie G, Cameron JI. Towards a universal model of family centered care: a scoping review. *BMC Health Serv Res* 2019;19:564. <https://doi.org/10.1186/s12913-019-4394-5>.
  98. Ditto PH, Jacobson JA, Smucker WD, Danks JH, Fagerlin A. Context changes choices: a prospective study of the effects of hospitalization on life-sustaining treatment preferences. *Med Decis Making* 2006;26:313–22. <https://doi.org/10.1177/0272989X06290494>.
  99. Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ* 2010;340. c1345. <https://doi.org/10.1136/bmj.c1345>.
  100. Cook D, Swinton M, Toledo F, Clarke F, Rose T, Hand-Breckenridge T, Boyle A, Woods A, Zytaruk N, Heels-Ansdell D, et al. Personalizing death in the intensive care unit: the 3 Wishes Project: a mixed-methods study. *Ann Intern Med* 2015;163:271–9. <https://doi.org/10.7326/M15-0502>.
  101. Reid JC, Dennis B, Hoard N, Clarke F, Hanmiah R, Vegas DB, Boyle A, Toledo F, Rudkowski JC, Soth M, et al. Enhancing end of life care on general internal medical wards: the 3 Wishes Project. *BMC Palliat Care* 2023;22:11. <https://doi.org/10.1186/s12904-023-01133-4>.
  102. Douma MJ, Graham TAD, Ali S, Dainty KN, Bone A, Smith KE, Dennet L, Brindley PG, Kroll T, Frazer K. What are the care needs of families experiencing cardiac arrest?: a survivor and family led scoping review. *Resuscitation* 2021;168:119–41. <https://doi.org/10.1016/j.resuscitation.2021.09.019>.
  103. Carlsson N, Arestedt K, Alvariza A, Axelsson L, Bremer A. Factors associated with symptoms of prolonged grief and psychological distress among bereaved family members of persons who died from sudden cardiac arrest. *J Cardiovasc Nurs* 2022;38:454–61. <https://doi.org/10.1097/JCN.0000000000000937>.
  104. Harrod M, Kamphuis LA, Hauschildt K, Seigworth C, Korpela PR, Rouse M, Vincent BM, Nallamothu BK, Iwashyna TJ. Getting better or getting by?: a qualitative study of in-hospital cardiac arrest survivors long-term recovery experiences. *SSM Qual Res Health*. 2021;1. 100002. <https://doi.org/10.1016/j.ssmqr.2021.100002>.
  105. Lijja G. Follow-up of cardiac arrest survivors: why, how, and when? A practical approach. *Semin Neurol* 2017;37:88–93. <https://doi.org/10.1055/s-0036-1593859>.
  106. McMeekin DE, Hickman Jr RL, Douglas SL, Kelley CG. Stress and coping of critical care nurses after unsuccessful cardiopulmonary resuscitation. *Am J Crit Care* 2017;26:128–35. <https://doi.org/10.4037/aicc2017916>.
  107. Kozelj A, Sikic Pogacar M, Fijan S, Strauss M, Postuvan V, Strnad M. Exploring the feelings of nurses during resuscitation—a cross-sectional study. *Healthcare (Basel)*. 2021;10:5. <https://doi.org/10.3390/healthcare10010005>.
  108. Spencer SA, Nolan JP, Osborn M, Georgiou A. The presence of psychological trauma symptoms in resuscitation providers and an exploration of debriefing practices. *Resuscitation* 2019;142:175–81. <https://doi.org/10.1016/j.resuscitation.2019.06.280>.
  109. Cole FL, Slocumb EM, Muldoon MJ. A measure of critical care nurses' post-code stress. *J Adv Nurs* 2001;34:281–8. <https://doi.org/10.1046/j.1365-2648.2001.01756.x>.
  110. Hodkinson A, Zhou A, Johnson J, Geraghty K, Riley R, Panagopoulou E, Chew-Graham CA, Peters D, Esmail A, Panagioti M. Associations of physician burnout with career engagement and quality of patient care: systematic review and meta-analysis. *BMJ* 2022;378:e070442.
  111. Green AA, Kinchen EV. The effects of mindfulness meditation on stress and burnout in nurses. *J Holist Nurs* 2021;39:356–68. <https://doi.org/10.1177/08980101211015818>.
  112. Copeland D, Liska H. Implementation of a post-code pause: extending post-event debriefing to include silence. *J Trauma Nurs* 2016;23:58–64. <https://doi.org/10.1097/JTN.0000000000000187>.
  113. Joe MB, Cusano A, Leckie J, Czuczman N, Exner K, Yong H, Ruzycy S, Lithgow K. Mentorship programs in residency: a scoping review. *J Grad Med Educ* 2023;15:190–200. [https://doi.org/10.4300/JGME-D-22-00415\\_1](https://doi.org/10.4300/JGME-D-22-00415_1).
  114. Fleming K. Reducing staff turnover through the implementation of a peer mentoring program. *Nurs Manage* 2023;54:32–9. <https://doi.org/10.1097/01-NUMA.0000905016.75550.3f>.
  115. Bonifacino E, Ufomata EO, Farkas AH, Turner R, Corbelli JA. Mentorship of underrepresented physicians and trainees in academic medicine: a systematic review. *J Gen Intern Med* 2021;36:1023–34. <https://doi.org/10.1007/s11606-020-06478-7>.
  116. Pethrick H, Nowell L, Paolucci EO, Lorenzetti L, Jacobsen M, Clancy T, Lorenzetti DL. Peer mentoring in medical residency education: a systematic review. *Can Med Educ J*. 2020;11: e128–37. <https://doi.org/10.36834/cmiej.68751>.
  117. Albott CS, Wozniak JR, McGlinch BP, Wall MH, Gold BS, Vinogradov S. Battle buddies: rapid deployment of a psychological resilience intervention for health care workers during the COVID-19 pandemic. *Anesth Analg* 2020;131:43–54. <https://doi.org/10.1213/ANE.0000000000004912>.
  118. Greif R, Bhanji F, Bigham BL, Bray J, Breckwoldt J, Cheng A, Duff JP, Gilfoyle E, Hsieh MJ, Iwami T, et al. Education, Implementation, and Teams Collaborators. Education, implementation, and teams: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation* 2020;142(16\_suppl\_1):S222–83. <https://doi.org/10.1161/CIR.0000000000000896>.
  119. Kolbe M, Schmutz S, Seelandt JC, Eppich WJ, Schmutz JB. Team debriefings in healthcare: aligning intention and impact. *BMJ* 2021;374. n2042. <https://doi.org/10.1136/bmj.n2042>.
  120. Sweberg T, Sen AI, Mullan PC, Cheng A, Knight L, Del Castillo J, Ikeyama T, Seshadri R, Hazinski MF, Raymond T, et al. Pediatric Resuscitation Quality (pediRES-Q) Collaborative Investigators. Description of hot debriefings after in-hospital cardiac arrests in an international pediatric quality improvement collaborative.

- Resuscitation 2018;128:181–7. <https://doi.org/10.1016/j.resuscitation.2018.05.015>.
121. Harder N, Lemoine J, Harwood R. Psychological outcomes of debriefing healthcare providers who experience expected and unexpected patient death in clinical or simulation experiences: a scoping review. *J Clin Nurs* 2020;29:330–46. <https://doi.org/10.1111/jocn.15085>.
  122. Dewan M, Parsons A, Tegtmeyer K, Wenger J, Niles D, Raymond T, Cheng A, Skellett S, Roberts J, Jani P, et al. Contextual factors affecting implementation of in-hospital pediatric CPR quality improvement interventions in a resuscitation collaborative. *Pediatr Qual Saf* 2021;6:e455. <https://doi.org/10.1097/pq9.0000000000000455>.
  123. Schnaubelt S, Garg R, Atiq H, Baig N, Bernardino M, Bigham B, Dickson S, Geduld H, Al-Hilali Z, Karki S, et al. Cardiopulmonary Resuscitation in Low-Resource Settings Group. Cardiopulmonary resuscitation in low-resource settings: a statement by the International Liaison Committee on Resuscitation, supported by the AFEM, EUSEM, IFEM, and IFRC. *Lancet Glob Health* 2023;11: e1444–53. [https://doi.org/10.1016/S2214-109X\(23\)00302-9](https://doi.org/10.1016/S2214-109X(23)00302-9).